

1/PRTS

09/719219

528 Rec CT/PTO 11 DEC 2000

WO 99/65872

PCT/FR99/01389

**DIARYLSELENIDE COMPOUNDS AND USE THEREOF IN HUMAN OR
VETERINARY MEDICINE AND IN COSMETICS**

The invention relates, as novel and useful industrial products, to diarylselenide compounds. The
5 invention also relates to the use of these novel compounds in pharmaceutical compositions intended for use in human or veterinary medicine, or alternatively in cosmetic compositions.


The compounds according to the invention have
10 pronounced activity in the fields of cell proliferation and differentiation and find applications more particularly in the topical and systemic treatment of dermatological complaints associated with a
keratinization disorder, dermatological (or other)
15 complaints with an inflammatory and/or immunoallergic component, and dermal or epidermal proliferations, whether benign or malignant. These compounds can also be used in the treatment of degenerative diseases of connective tissue, to combat ageing of the skin,
20 whether light-induced or chronological, and to treat cicatrization disorders. They moreover find an application in the ophthalmological field, in particular in the treatment of corneopathies.

The compounds according to the invention can
25 also be used in cosmetic compositions for body and hair hygiene.

The present invention relates to compounds

09719219-032801

which can be represented by the general formula (I) below:



(I)


in which:

5 - R_1 represents:

- (i) a $-CH_3$ radical,
- (ii) a radical $-CH_2-O-R_5$,
- (iii) a radical $-COR_6$,

R_5 and R_6 having the meanings given below,

10 - Ar represents a radical chosen from the radicals of formulae (a)-(e) below:



R_7 having the meaning given below,

- R_2 and R_3 , which may be identical or different,

15 independently represent a radical chosen from:

- (i) a hydrogen atom,
- (ii) a radical chosen from tert-butyl, 1-methylcyclohexyl and 1-adamantyl radicals,
- (iii) a radical $-OR_8$, R_8 having the meaning given below,
- (iv) a polyether radical,

20

it being understood that at least one of the radicals R_2 or R_3 represents a radical (ii),

- R_2 and R_3 taken together can form, with the adjacent aromatic ring, a 5- or 6-membered saturated ring

5 optionally substituted with methyl groups and/or optionally interrupted with an oxygen or sulphur atom,

- R_4 represents a hydrogen atom, a halogen atom, a lower alkyl radical, a radical OR_9 , a polyether radical or a radical COR_{10} ,

10 R_9 and R_{10} having the meanings given below,

- R_5 represents a hydrogen atom, a lower alkyl radical or a radical COR_{11} ,


R_{11} having the meaning given below,

- R_6 represents a radical chosen from:

- 15 (i) a hydrogen atom,
(ii) a lower alkyl radical,
(iii) a radical OR_{12} ,

R_{12} having the meaning given below,

(iv) a radical of formula




20

R' and R'' having the meanings given below,

- R_7 represents a hydrogen atom, a halogen atom, a lower alkyl radical, a nitro radical, a radical OR_{13} , a polyether radical or a radical of the following

25 formula:

T082ED 6T26T60



R_{13} , R_{14} and R_{15} having the meanings given below,

- R_8 represents a hydrogen atom, a lower alkyl radical, an optionally substituted aryl radical, an optionally substituted aralkyl radical, a monohydroxyalkyl or polyhydroxyalkyl radical or a lower acyl radical,
- R_9 represents a hydrogen atom, a lower alkyl radical, an optionally substituted aryl radical, an optionally substituted aralkyl radical, a monohydroxyalkyl or polyhydroxyalkyl radical, a lower acyl radical, a radical $-(\text{CH}_2)_n\text{-COOR}_{16}$ or a radical $-(\text{CH}_2)_n\text{-X}$,
 n , R_{16} and X having the meanings given below,
- R_{10} and R_{11} , which may be identical or different, represent a lower alkyl radical,
- R_{12} represents a hydrogen atom, a lower alkyl radical, an optionally substituted aryl or aralkyl radical, a monohydroxyalkyl radical or a polyhydroxyalkyl radical,
- R' and R'' , which may be identical or different, represent a hydrogen atom, a lower alkyl radical, an optionally substituted aryl radical or an amino acid residue,
 or alternatively R' and R'' taken together can form, with the nitrogen atom, a heterocycle,
- R_{13} represents a hydrogen atom or a lower alkyl radical,

09719249-032801
TOPSECRET

- R_{14} and R_{15} , which may be identical or different, represent a hydrogen atom or a lower alkyl radical,
 - R_{16} represents a hydrogen atom or a lower alkyl radical,
- 5 - n represents an integer between 1 and 12 inclusive,
- X represents a halogen atom.

The invention is also directed towards the salts of the compounds of formula (I) when R_1 represents a carboxylic acid function, and the geometrical and
10 optical isomers of the said compounds of formula (I).

When the compounds according to the invention are in the form of salts, they are preferably salts of an alkali metal or alkaline-earth metal, or alternatively of zinc or of an organic amine.

15 According to the present invention, the expression "lower alkyl radical" means a radical containing from 1 to 6 carbon atoms, and preferably methyl, ethyl, isopropyl, butyl and tert-butyl radicals.

20 The expression "monohydroxyalkyl radical" means a radical containing from 1 to 6 carbon atoms, in particular a 2-hydroxyethyl, 2-hydroxypropyl or 3-hydroxypropyl radical, it being possible for the monohydroxyalkyl radical to be protected in the form of
25 acetyl or tert-butyldimethylsilyl.

The expression "polyhydroxyalkyl radical" means a radical containing from 2 to 6 carbon atoms and from 2 to 5 hydroxyl groups, such as, in particular,

05719619-032804

2,3-dihydroxypropyl, 2,3,4-trihydroxybutyl,
2,3,4,5-tetrahydroxypentyl radicals or a
pentaerythritol residue, it being possible for the
hydroxyl groups to be protected in the form of acetyls
5 or tert-butyldimethylsilyls.

The expression "optionally substituted aryl
radical" means a phenyl radical optionally substituted
with at least one halogen atom, a hydroxyl optionally
protected in the form of an ether or acetate function,
10 a nitro function or an amino function optionally
substituted with an alkyl or acetyl group.

The expression "optionally substituted
aralkyl radical" means a benzyl radical or a phenethyl
radical optionally substituted with at least one
15 halogen atom, a hydroxyl radical optionally protected
in the form of an ether or acetate function, a nitro
function or an amino function optionally substituted
with an alkyl or acetyl group.

The expression "lower acyl radical" means a
20 radical containing from 1 to 4 carbon atoms, and
preferably an acetyl or propionyl radical.

The expression "amino acid residue" means a
residue derived, for example, from one of the 20 amino
acids of L or D configuration which constitute
25 mammalian proteins.

The term "heterocycle" preferably means a
piperidino, morpholino, pyrrolidino or piperazino
radical, optionally substituted in position 4 with a

03719219 032801
TOB220 51261250

C₁-C₆ alkyl radical or with a mono- or polyhydroxyalkyl radical as defined above.

The expression "polyether radical" means a radical containing from 1 to 6 carbon atoms and from 1 to 3 oxygen or sulphur atoms, such as methoxymethyl ether, methoxyethoxymethyl ether or methylthiomethyl ether radicals.

The expression "halogen atom" preferably means a fluorine, chlorine or bromine atom.

10 According to the present invention, the compounds of formula (I) that are more particularly preferred are those for which at least one, and preferably all, of the conditions below are satisfied:

- R₁ represents a radical COR₆
- 15 - Ar represents a radical of formula (a) or (b)
- R₂ or R₃ represents an adamantyl radical or R₂ and R₃ taken together form, with the adjacent aromatic ring, a 5- or 6-membered saturated ring optionally substituted with methyl groups and/or optionally
- 20 interrupted with an oxygen or sulphur atom.

Among the compounds of formula (I) above falling within the context of the present invention, mention may be made in particular of the following:

ethyl 4-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate,

25 4-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid,

ethyl 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-

- 2-naphthylselanyl)nicotinate,
 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)nicotinic acid,
 ethyl 6-(5,5,8,8-tetramethyl-3-propoxy-5,6,7,8-
 5 tetrahydro-2-naphthylselanyl)nicotinate,
 6-(5,5,8,8-tetramethyl-3-propoxy-5,6,7,8-tetrahydro-
 2-naphthylselanyl)nicotinic acid,
 3-(4-tert-butylphenylselanyl)benzoic acid,
 6-(4-tert-butylphenylselanyl)nicotinic acid,
 10 4-(4-tert-butylphenylselanyl)benzoic acid,
 4-(4,4-dimethylthiochroman-8-ylselanyl)benzoic acid,
 3-(4,4-dimethylthiochroman-8-ylselanyl)benzoic acid,
 6-(4,4-dimethylthiochroman-8-ylselanyl)nicotinic acid,
 4-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthyl-
 15 selanyl)benzoic acid,
 3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)benzoic acid,
 6-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)nicotinic acid,
 20 4-[5-adamantan-1-yl-4-(2-methoxyethoxymethoxy)-
 2-methylphenylselanyl]benzoic acid,
 3-[5-adamantan-1-yl-4-(2-methoxyethoxymethoxy)-
 2-methylphenylselanyl]benzoic acid,
 6-(4-methoxyethoxymethoxy-5,5,8,8-tetramethyl-
 25 5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid,
 3-(4-methoxyethoxymethoxy-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid,
 4-(4-methoxyethoxymethoxy-5,5,8,8-tetramethyl-

09719219-032804
 TOPSECRET

- 5,6,7,8-tetrahydro-2-naphthylselanyl)-3-methoxybenzoic acid,
- 3-(4-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-4-methoxybenzoic acid,
- 5 6-(4-methoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid,
- 6-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid,
- 10 2-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid,
- 4-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid,
- 3-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid,
- 15 6-(3,5-di-tert-butyl-2-methoxymethoxyphenylselanyl)-nicotinic acid,
- 2-(3,5-di-tert-butyl-2-methoxymethoxyphenylselanyl)-nicotinic acid,
- 20 4-(3,5-di-tert-butyl-2-methoxymethoxyphenylselanyl)-benzoic acid,
- 3-(3,5-di-tert-butyl-2-methoxymethoxyphenylselanyl)-benzoic acid,
- 6-[4-adamantan-1-yl-3-benzyloxyphenylselanyl]nicotinic acid,
- 25 6-(3,5-di-tert-butyl-2-benzyloxyphenylselanyl)nicotinic acid,
- 3-methoxy-4-(4-benzyloxy-5,6,7,8-tetrahydro-

F09250-5T26T/60

- 5,5,8,8-tetramethyl-2-naphthylselanyl)benzoic acid,
 4-(4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
 2-naphthylselanyl)benzoic acid,
 6-(4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
 5 2-naphthylselanyl)nicotinic acid,
 3-methoxy-4-(3-benzyloxy-5,6,7,8-tetrahydro-
 5,5,8,8-tetramethyl-2-naphthylselanyl)benzoic acid,
 6-(3-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
 2-naphthylselanyl)nicotinic acid,
 10 4-(3-hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
 2-naphthylselanyl)-3-methoxybenzoic acid,
 6-(3-hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
 2-naphthylselanyl)nicotinic acid,
 4-(5-adamantan-1-yl-4-benzyloxy-2-methylphenylselanyl)-
 15 benzoic acid,
 6-[3-(5-hydroxypentyloxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic acid,
 ethyl 4-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)benzoate,
 20 ethyl 4-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate,
 ethyl 4-(3-hydroxy-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate,
 4-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 25 2-naphthylselanyl)benzoic acid,
 ethyl 6-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate,
 ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-

09719219-0322004
 TOPSECRET

- 5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate,
 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)nicotinic acid,
 ethyl 6-[3-(3-ethoxycarbonylpropoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate,
 6-[3-(3-carboxypropoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic acid,
 ethyl 4-[3-(3-ethoxycarbonylpropoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]benzoate,
 4-[3-(3-carboxypropoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]benzoic acid,
 ethyl 4-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetra-
 methyl-5,6,7,8-tetrahydro-2-naphthylselanyl]benzoate,
 4-[3-(7-carboxyheptyloxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]benzoic acid,
 ethyl 6-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetra-
 methyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate,
 6-[3-(7-carboxyheptyloxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic acid,
 ethyl 6-[3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate,
 6-[3-(2-hydroxyethoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic acid,
 ethyl 4-[3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]benzoate,
 4-[3-(2-hydroxyethoxy)-5,5,8,8-tetramethyl-
 5,6,7,8-tetrahydro-2-naphthylselanyl]benzoic acid,
 6-(3-adamantan-1-yl-4-methoxyphenylselanyl)nicotinic

T09250-67257-60

acid,

[6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-

2-naphthylselanyl)-3-pyridyl]methanol,

N-ethyl-6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-

5 2-naphthylselanyl)nicotinamide,

morpholin-4-yl-[6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetra-

hydro-2-naphthylselanyl)-3-pyridyl]methanone,

N-(4-hydroxyphenyl)-6-(3,5,5,8,8-pentamethyl-

5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinamide,

10 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-

2-naphthylselanyl)pyridine-3-carbaldehyde.

Subjects of the present invention are also processes for preparing the compounds of formula (I), in particular according to the reaction scheme given in

15 Figure 1.

The derivatives of formula (I) can be obtained (Fig. 1) by a sequence of reactions comprising the action of a lithiated base such as tBuLi on the product (2) in a solvent such as THF, followed by
20 addition of selenium and the formation of the dimer by oxidation in basic medium (EtOH, NaOH). The product (3) obtained is subjected to the action of sodium borohydride in a solvent such as ethanol and then coupled with an iodoaryl in the presence of a nickel
25 catalyst.

When R_1 represents a COOH radical, the compounds are prepared by protecting R_1 with a protecting group of alkyl type. Saponification of the

09719249-032804

ester function in the presence of a base, such as sodium hydroxide or lithium hydroxide in an alcoholic solvent or in THF, gives the corresponding acids.

When R_1 represents an alcohol radical, the
5 compounds can be obtained from the acid by reduction in
the presence of hydride such as boron hydride. The
alcohol can be etherified according to the conventional
methods.

When R_1 represents an aldehyde radical, the
10 compounds can be obtained by oxidation of the
corresponding alcohols by the action of manganese oxide
or pyridinium dichromate.

When R_1 represents an amide radical, the compounds can be obtained by converting the acid into the acid chloride and then by reaction with a suitable amine.

These compounds bind to RXR receptors, some having agonist activity, others having antagonist activity.

20 The binding and transactivation properties as
RXR receptor agonists can be determined by methods
known in the art, such as, for example: Levin et al.,
Nature 1992, **355**, 359-61; Allenby et al., Proc. Natl.
Acad. Sci., 1993, **90**, 30-4.

25 The RXR-agonist activity can also be
determined by the test as described in French patent
application No. 95/07301 filed on 19 June 1995 by the
Applicant. This test comprises the following steps:

(i) a sufficient amount of a compound which is an active ligand of at least one receptor of the steroidal/thyroid nuclear receptor superfamily, other than an RXR-receptor-specific ligand which can

5 heterodimerize with the RXRs such as an RAR-agonist molecule, is applied topically to an area of skin of a mammal, (ii) a molecule capable of presenting RXR-agonist activity is administered systemically or topically to this same area of mammalian skin before,

10 during or after step (i), and (iii) the response on the area of mammal's skin thus treated is evaluated. Thus, the response to a topical application to a mammal's ear of an RAR-agonist molecule which corresponds to an increase in the thickness of this ear can be increased

15 by administering an RXR-receptor-agonist molecule systemically or topically.

The RXR α -antagonist activity can be evaluated in the transactivation test by determination of the dose (IC₅₀) which gives 50% inhibition of the

20 transactivating activity of an RXR α -selective agonist: 6-[1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)cyclopropyl]nicotinic acid (CD 3127) according to the following procedure:

HeLa cells are co-transfected with an

25 expression vector coding for RXR α (p565-RXR α) and a reporter plasmid containing the response element 1/2 CRBP II cloned upstream of the thymidine kinase heterologous promoter and of the chloramphenicolm-

09719219-032801
103250-6726760

acetyl-transferase (CAT) reporter gene. Eighteen hours after co-transfection, the cells are treated with a fixed concentration of CD 3127 and increasing concentrations of the molecule to be evaluated. After treatment for twenty-four hours, the CAT activity is assayed by ELISA. The fixed concentration of CD3127 used is 10^{-8} M and corresponds to its EC_{50} .

A subject of the present invention is thus, as a medicinal product, the compounds of formula (I) as defined above.

The compounds according to the invention are particularly suitable in the following fields of treatment:

1) for treating dermatological complaints associated with a keratinization disorder which has a bearing on differentiation and on proliferation, in particular for treating common acne, comedones, polymorphonuclear leukocytes, rosacea, nodulocystic acne, acne conglobata, senile acne, secondary acnes such as solar, medication-related or occupational acne,

2) for treating other types of keratinization disorder, in particular ichthyosis, ichthyosiform states, Darier's disease, palmoplantar keratoderma, leucoplasias and leucoplasiform states, and cutaneous or mucous (buccal) lichen,

3) for treating other dermatological complaints associated with a keratinization disorder with an inflammatory and/or immunoallergic component and, in

particular, all forms of psoriasis, whether it is cutaneous, mucous or unguual psoriasis and even psoriatic rheumatism, or alternatively cutaneous atopy, such as eczema or respiratory atopy or alternatively gingival hypertrophy; the compounds can also be used in certain inflammatory complaints which have no keratinization disorder,

4) for treating all dermal or epidermal proliferations, whether benign or malignant and whether they are of viral origin or otherwise, such as common warts, flat warts and verruciform epidermodysplasia, oral or florid papillomatoses and proliferations which may be induced by ultraviolet radiation, in particular in the case of basocellular and spinocellular epithelioma,

5) for treating other dermatological disorders such as bullosis and collagen diseases,

6) for treating certain ophthalmological disorders, in particular corneopathies,

7) for repairing or combating ageing of the skin, whether this is light-induced or chronological ageing, or for reducing actinic keratoses and pigmentations, or any pathologies associated with chronological or actinic ageing,

8) for preventing or curing the stigmata of epidermal and/or dermal atrophy induced by local or systemic corticosteroids, or any other form of cutaneous atrophy,

09719219-032801

9) for preventing or treating cicatrization disorders or for preventing or repairing stretchmarks, or alternatively for promoting cicatrization,

10) for combating disorders of sebaceous
5 functioning such as the hyperseborrhoea of acne or simple seborrhoea,

11) in the treatment or prevention of cancerous or precancerous states, more particularly promyelocyte leukaemias,

10 12) in the treatment of inflammatory complaints such as arthritis,

13) in the treatment of any general or skin complaint of viral origin,

14) in the prevention or treatment of alopecia,

15 15) in the treatment of dermatological or general complaints having an immunological component,

16) in the treatment of complaints of the cardiovascular system such as arteriosclerosis, hypertension, non-insulin-dependent diabetes and
20 obesity,

17) in the treatment of skin disorders due to an exposure to U.V. radiation.

In the therapeutic fields mentioned above, the compounds according to the invention may be
25 employed advantageously in combination with other compounds of retinoid-type activity, with D vitamins or derivatives thereof, with corticosteroids, with anti-free-radical agents, α -hydroxy or α -keto acids or

094849-0330
T0330-5726260

derivatives thereof, or alternatively with ion-channel blockers. The expression "D vitamins or derivatives thereof" means, for example, vitamin D₂ or D₃ derivatives and in particular 1,25-dihydroxyvitamin D₃.

5 The expression "anti-free-radical agents" means, for example, α -tocopherol, superoxide dismutase, ubiquinol or certain metal-chelating agents. The expression " α -hydroxy or α -keto acids or derivatives thereof" means, for example, lactic, malic, citric, glycolic,
10 mandelic, tartaric, glyceric or ascorbic acid salicylic acid derivatives, or the salts, amides or esters thereof. Lastly, the term "ion-channel blockers" means, for example, Minoxidil (2,4-diamino-6-
15 piperidinopyrimidine 3-oxide) and derivatives thereof.

A subject of the present invention is also medicinal compositions containing at least one compound of formula (I) as defined above, one of the optical or geometrical isomers thereof or one of the salts thereof.

20 A subject of the present invention is thus a novel medicinal composition intended in particular for treating the abovementioned complaints, and which is characterized in that it comprises, in a pharmaceutically acceptable support which is compatible
25 with the mode of administration selected for this composition, at least one compound of formula (I), one of the optical or geometrical isomers thereof or one of the salts thereof.

The compounds according to the invention may be administered enterally, parenterally, topically or ocularly.

Via the enteral route, the medicinal products
5 may be in the form of tablets, gelatin capsules, sugar-coated tablets, syrups, suspensions, solutions, powders, granules, emulsions, microspheres or nanospheres or polymeric or lipid vesicles which enable controlled release. Via the parenteral route, the
10 compositions may be in the form of solutions or suspensions for infusion or for injection.

The compounds according to the invention are generally administered at a daily dose of about 0.01 mg/kg to 100 mg/kg of body weight taken in 1 to 3
15 doses.

Via the topical route, the pharmaceutical compositions based on compounds according to the invention are more particularly intended for the treatment of the skin and the mucosae and may thus be
20 in the form of ointments, creams, milks, salves, powders, impregnated pads, solutions, gels, sprays, lotions or suspensions. They may also be in the form of microspheres or nanospheres or polymeric or lipid vesicles or polymeric patches and hydrogels which
25 enable controlled release. These topical-route compositions may either be in anhydrous form or in aqueous form, depending on the clinical indication.

09749249.0322001
T0322001.6T26T460

Via the ocular route, they are mainly
eyedrops.

These compositions for topical or ocular use
contain at least one compound of formula (I) as defined
5 above, or one of the optical or geometrical isomers
thereof or alternatively one of the salts thereof, at a
concentration preferably of between 0.001% and 5% by
weight relative to the total weight of the composition.

The compounds of formula (I) according to the
10 invention also find an application in the cosmetic
field, in particular in body and hair hygiene and
especially for treating skin types with a tendency
towards acne, for promoting the regrowth of the hair,
for combating hair loss, for combating the greasy
15 appearance of the skin or the hair, in protection
against the harmful effects of the sun or in the
treatment of physiologically dry skin types, and for
preventing and/or combating light-induced or
chronological ageing.

20 In the cosmetic field, the compounds
according to the invention can moreover be employed
advantageously in combination with other compounds of
retinoid-type activity, with D vitamins or derivatives
thereof, with corticosteroids, with anti-free-radical
25 agents, α -hydroxy or α -keto acids or derivatives
thereof, or alternatively with ion-channel blockers,
all of these various products being as defined above.

09719219-032801

09719219-032301

The present invention is thus also directed towards a cosmetic composition which is characterized in that it comprises, in a cosmetically acceptable support which is suitable for topical application, at least one compound of formula (I) as defined above or one of the optical or geometrical isomers thereof or one of the salts thereof, it being possible for this cosmetic composition to be, in particular, in the form of a cream, a milk, a lotion, a gel, microspheres or nanospheres or polymeric or lipid vesicles, a soap or a shampoo.

The concentration of compound of formula (I) in the cosmetic compositions according to the invention is advantageously between 0.001% and 3% by weight relative to the composition as a whole.

The medicinal and cosmetic compositions according to the invention can also contain inert additives or even pharmacodynamically or cosmetically active additives or combinations of these additives and, in particular: wetting agents; depigmenting agents such as hydroquinone, azelaic acid, caffeic acid or kojic acid; emollients; moisturizing agents such as glycerol, PEG 400, thiamorpholinone and derivatives thereof, or urea; anti-seborrhoea or anti-acne agents such as S-carboxymethylcysteine, S-benzylcysteamine, the salts and the derivatives thereof, or benzoyl peroxide; antibiotics such as erythromycin and esters thereof, neomycin, clindamycin and esters thereof, and

The compositions according to the invention may also contain flavour-enhancing agents, preserving agents such as para-hydroxybenzoic acid esters, stabilizing agents, moisture regulators, pH regulators, osmotic pressure modifiers, emulsifying agents, UV-A and UV-B screening agents, and antioxidants such as α -tocopherol, butylhydroxyanisole or butylhydroxytoluene.

Several examples for obtaining active compounds of formula (I) according to the invention, as well as various concrete formulations based on such compounds, will now be given for illustrative purposes and with no limiting nature.

A. EXAMPLES OF COMPOUNDS**EXAMPLE 1:**

Ethyl 4-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)benzoate

- 5 (a) 5,6,7,8-Tetrahydro-3,5,5,8,8-pentamethylnaphthalene-2-diselenide

1.7 M tert-butyllithium in pentane
(37.4 mmol, 22 ml) is added to a solution of 2-bromo-5,6,7,8-tetrahydro-3,5,5,8,8-pentamethylnaphthalene
10 (4.4 g, 15.8 mmol) in THF (100 ml) at -78°C over 10 min. The mixture is stirred at 0°C for 30 min. Selenium (1.33 g, 16.8 mmol) is added in 2 portions. The mixture is stirred at 0°C for 15 min and then at room temperature for 30 min. 1N HCl solution (40 ml) is
15 added and the reaction mixture is then treated with ethyl ether. The organic phase is washed twice with water, dried over anhydrous magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C. 10 ml of ethanol and 50 mg of sodium hydroxide
20 are added to the oil obtained. The mixture is stirred vigorously for a few minutes in air (until the product has all precipitated) and is then concentrated on a rotary evaporator under vacuum at 40°C. The solid obtained is filtered through silica (eluting with
25 heptane) and then crystallized from an ethanol/ether mixture. Yellow solid. Mass: 3.26 g. Yield: 74%.
m.p.: 126°C.
1H NMR (CDCl₃): 1.14 (6H, s), 1.23 (6H, s), 1.61 (4H,

09719219-032801
TOP SECRET 676760

s), 2.35 (3H, s), 7.05 (1H Ar, s), 7.55 (1H Ar, s).

(b) Ethyl 4-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate

A solution of 5,6,7,8-tetrahydro-3,5,5,8,8-pentamethylnaphthalene-2-diselenide (500 g, 0.89 mmol) and sodium borohydride (68 mg, 1.8 mmol) in 5 ml of ethanol is stirred for 1 hour at room temperature. Ethyl iodobenzoate (440 mg, 1.6 mmol) and bis(bipyridine)nickel(II) bromide (10 mg, 0.016 mmol) (Organometallics 1985, 4, 657-661) are then added. The solution is refluxed for 5 minutes. At room temperature, it is diluted with ethyl ether. The organic phase is washed with water, dried over anhydrous magnesium sulphate and then concentrated. The residue is purified by fast plug (eluent: heptane and then ethyl ether).

White solid. Mass: 495 mg. Yield: 72%. m.p.: 104°C.

¹H NMR (CDCl₃): 1.22 (6H, s), 1.29 (6H, s), 1.33-1.39 (3H, t), 1.67 (4H, s), 2.32 (3H, s), 4.29-4.38 (2H, q), 7.21-7.26 (3H, c), 7.51 (1H, s), 7.84-7.87 (2H, d).

EXAMPLE 2:

4-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid

Sodium hydroxide (450 mg, 11.25 mmol) is added to a solution of ethyl 4-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate (450 mg, 1.04 mmol) in a mixture of 10 ml of THF, 1 ml of methanol and 1 ml of water. The reaction medium is

09719219-032801

refluxed for 12 h. It is then poured into an ethyl ether/water mixture, acidified to pH 1 with concentrated hydrochloric acid solution and extracted with ethyl ether. After separation of the phases by settling, the organic phase is washed twice with water, dried over anhydrous magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C.

White powder. Mass: 371 mg. Yield: 88%. m.p.: 249°C.

10 ¹H NMR (CDCl₃): 1.21 (6H, s), 1.29 (6H, s), 1.67 (4H, s), 2.32 (3H, s), 7.21-7.24 (2H, d, J=6.9 Hz), 7.38 (1H, s), 7.48 (1H, s), 7.85-7.88 (2H, d, J=8.35 Hz).

EXAMPLE 3:

15 **Ethyl 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate**

In a manner similar to that of Example 1(b), by reaction of 750 mg (1.33 mmol) of diselenide in 15 ml of ethanol with 102 mg (2.7 mmol) of sodium borohydride, 665 mg (2.4 mmol) of ethyl 6-iodonicotinate and 15 mg (0.024 mmol) of bis(bipyridine)nickel(II) bromide, 779 mg (75%) of the expected derivative are obtained in the form of a white solid. m.p.: 117°C.

25 ¹H NMR (CDCl₃): 1.25 (6H, s), 1.31 (6H, s), 1.34-1.40 (3H, t), 1.69 (4H, s), 2.37 (3H, s), 4.32-4.40 (2H, q), 6.83-6.87 (1H, d, J=8.3 Hz), 7.28 (1H, s), 7.65 (1H, s), 7.91-7.96 (1H, dd, J=6.10 Hz, J'=2.21 Hz), 8.99-9.00 (1H, d, J=2.14 Hz).

00719219-032201
T08220-6T26T260

EXAMPLE 4:**6-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid**

In a manner similar to that of Example 2, by
 5 reaction of 750 mg (1.74 mmol) of ethyl 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-nicotinate with 700 mg (17.5 mmol) of sodium hydroxide in a THF/methanol/water mixture, 625 mg (89%) of a white cottony product are obtained. m.p.: 258°C.

10 ¹H NMR (DMSO): 1.05 (6H, s), 1.11 (6H, s), 1.48 (4H, s), 2.14 (3H, s), 6.79-6.83 (1H, d, J=8.3 Hz), 7.24 (1H, s), 7.45 (1H, s), 7.83-7.88 (1H, dd, J=6.03 Hz, J'=2.3 Hz), 8.69-8.70 (1H, d, J=2.2 Hz), 13.12 (1H, s).

EXAMPLE 5:

15 **Ethyl 6-(5,5,8,8-tetramethyl-3-propoxy-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate**

(a) 1,1,4,4,-Tetramethyl-7-propoxy-1,2,3,4-tetrahydronaphthalene-6-diselenide

In a manner similar to that of Example 1(a),
 20 by reaction of 6 g (18.5 mmol) of 6-bromo-1,1,4,4-tetramethyl-7-propoxy-1,2,3,4-tetrahydronaphthalene with 1.7 M tert-butyllithium in pentane and selenium in 20 ml of THF, 3.2 g of the expected selenium derivative are obtained in the form of a yellow solid.

25 m.p.: 92-98°C.

¹H NMR (CDCl₃): 1.05-1.10 (6H, m), 1.25 (9H, m), 1.55-1.66 (4H, m), 1.86 (2H, sext), 3.98 (2H, t), 6.67 (1H, s), 7.42 (1H, s).

T08220"0326T260

(b) Ethyl 6-(5,5,8,8-tetramethyl-3-propoxy-5,6,7,8-tetrahydro-2-naphthylselenanyl)nicotinate

In a manner similar to that of Example 1(b), by reaction of 850 mg (1.31 mmol) of diselenide in 5 85 ml of ethanol with 120 mg (2.62 mmol) of sodium borohydride, 581 mg (2.1 mmol) of ethyl 6-iodonicotinate and 20 mg (0.032 mmol) of bis(bipyridine)nickel(II) bromide, 610 mg (61%) of the expected compound are obtained in the form of white 10 crystals. m.p.: 110-112°C.

¹H NMR (CDCl₃): 0.81-0.87 (3H, t), 1.24 (6H, s), 1.31 (6H, s), 1.35-1.41 (3H, t), 1.57-1.65 (2H, m), 1.69 (4H, s), 3.87-3.92 (2H, t), 4.32-4.41 (2H, q), 6.66 (1H, s), 7.00-7.03 (1H, d, J=8.3 Hz), 7.59 (1H, s), 15 7.91-7.95 (1H, dd, J=6.2 Hz, J'=2.1 Hz), 8.98-8.99 (1H, d, J=1.7 Hz).

EXAMPLE 6:

6-(5,5,8,8-Tetramethyl-3-propoxy-5,6,7,8-tetrahydro-2-naphthylselenanyl)nicotinic acid

20 In a manner similar to that of Example 2, by reaction of 485 mg (1.02 mmol) of ethyl 6-(5,5,8,8-tetramethyl-3-propoxy-5,6,7,8-tetrahydro-2-naphthylselenanyl)nicotinate with 385 mg (9.6 mmol) of sodium hydroxide in ethanol (20 ml), 444 mg (97%) of a 25 white solid are obtained. m.p.: 220°C.

EXAMPLE 7:

3-(4-tert-Butylphenylselenanyl)benzoic acid

A mixture of 4-tert-butylphenyl diselenide

(0.3 mmol), 480 mg of borohydride polymer supported on Amberlyst IRA 400 resin at 2.5 mmol/g (Aldrich), bis(bipyridine)nickel(II) dibromide (5 mg) (Organometallics 1985, 4, 657-661) and ethyl

5 3-iodobenzoate (0.4 mmol) is heated for 12 h at 67°C.

The mixture is filtered and the solution is concentrated. The solid obtained is purified on an SPE cartridge packed with silica gel. The fractions containing the expected product are combined and

10 concentrated under vacuum. The ester is saponified in a mixture of 2.5 ml of THF, 2.5 ml of ethyl alcohol and 0.5 ml of aqueous 33% sodium hydroxide solution. The reaction medium is acidified with HCl solution, extracted with ethyl ether, dried over magnesium

15 sulphate and concentrated to give the expected product.

¹H NMR/CDCl₃: 1.32 (s, 9H); 7.32 to 7.38 (m, 3H); 7.46 (d, 2H); 7.61 (d, 1H); 7.95 (d, 1H); 8.19 (d, 1H).

EXAMPLE 8:

6-(4-*tert*-Butylphenylselenalyl)nicotinic acid

20 The product is obtained in a manner similar to that of Example 7, starting with 4-*tert*-butylphenyl diselenide and ethyl 6-iodonicotinate.

¹H NMR/CDCl₃: 1.36 (s, 9H); 7.02 (d, 1H); 7.45 (d, 2H); 7.65 (d, 2H); 7.96 (d, 1H); 9.05 (d, 1H).

25 **EXAMPLE 9:**

4-(4-*tert*-Butylphenylselenalyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 4-*tert*-butylphenyl

T08220" 6T26T260

diselenide and ethyl 4-iodobenzoate.

^1H NMR/ CDCl_3 : 1.34 (s, 9H); 7.35 (d, 2H); 7.39 (d, 2H); 7.54 (d, 2H); 7.92 (d, 2H).

EXAMPLE 10

5 4-(4,4-Dimethylthiochroman-8-ylselenalyl)benzoic acid

(a) 2-Bromo-1-(3-methylbut-2-enylthio)benzene

19.30 g (102.0 mmol) of 2-bromothiophenol, 160 ml of DMF and 15.50 g (112.0 mmol) of potassium carbonate are introduced into a three-necked flask.

10 13 ml (112.0 mmol) of 1-bromo-3-methyl-2-butene are added dropwise and the mixture is stirred at room temperature for two hours. The reaction medium is poured into water and extracted with ethyl acetate, and the organic phase is separated out by settling, washed
15 with water, dried over magnesium sulphate and evaporated. 26.00 g (99%) of the expected compound are collected in the form of an orange-coloured oil.

^1H NMR (CDCl_3) δ 1.65 (s, 3H), 1.73 (s, 3H), 3.56 (d, 2H, $J=7.7$ Hz), 5.32 (td, 1H, $J=7.7/1.4$ Hz), 6.96 to
20 7.06 (m, 1H), 7.22 to 7.26 (m, 2H), 7.52 (d, 1H, $J=7.7$ Hz).

(b) 4,4-Dimethyl-8-bromothiochroman

26.00 g (102.0 mmol) of 2-bromo-1-(3-methylbut-2-enylthio)benzene, 180 ml of toluene
25 and 23.20 g (122.0 mmol) of para-toluenesulphonic acid are introduced into a three-necked flask. The reaction medium is refluxed for four hours and evaporated to dryness. The residue is taken up in aqueous sodium

09719219-032801

hydrogen carbonate solution and extracted with ethyl acetate, and the organic phase is separated out by settling, dried over magnesium sulphate and evaporated. The residue obtained is purified by chromatography on a column of silica, eluting with heptane. 20.00 g (76%) of the expected compound are collected in the form of an orange-coloured oil.

^1H NMR (CDCl_3) δ 1.33 (s, 6H), 1.94 (t, 2H, $J=6.0$ Hz), 3.04 (t, 2H, $J=6.1$ Hz), 6.89 (t, 1H, $J=7.9$ Hz), 7.34 (d, 2H, $J=7.9$ Hz).

(c) 4,4-Dimethylthiochroman-8-diselenide

One crystal of iodine, magnesium (208 mg, 8.56 mmol) and a few drops of a solution of 4,4-dimethyl-8-bromothiochroman (2 g, 7.78 mmol) in ethyl ether (15 ml) are heated until the organomagnesium reagent has been initiated. The rest of the solution is then added dropwise. The reaction medium is heated for 2 h and selenium (615 mg, 7.78 mmol) is then added at room temperature. Stirring is continued for 30 min and 1N HCl solution is then added. The reaction mixture is treated with ethyl ether. The organic phase is washed twice with water, dried over anhydrous magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C. Ethanol and sodium hydroxide are added to the oil obtained. The mixture is stirred vigorously for a few minutes and is then concentrated on a rotary evaporator under vacuum at 40°C.

The product is purified on a column of silica (20 dichloromethane/80 heptane).

White solid. Mass: 300 mg. Yield: 15%.

¹H NMR (CDCl₃): 1.33 (6H, s), 1.96 (2H, m), 3.09 (2H, m), 6.93 (1H Ar, t, J=7.8 Hz), 7.26 (1H Ar, dd, J=7.8 Hz, J=1.3 Hz), 7.47 (1H Ar, dd, J=7.8 Hz, J=1.3 Hz).

(d) 4-(4,4-Dimethylthiochroman-8-ylselenalyl)benzoic acid

10 The product is obtained in a manner similar to that of Example 7, starting with 4,4-dimethylthiochroman-8-diselenide and ethyl 4-iodobenzoate.

¹H NMR/CDCl₃: 1.36 (s, 6H); 1.95 (m, 2H), 2.99 (m, 2H), 6.99 (t, 1H), 7.31 to 7.46 (m, 4H); 7.91 (d, 2H).

15 **EXAMPLE 11:**

3-(4,4-Dimethylthiochroman-8-ylselenalyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 4,4-dimethylthiochroman-8-diselenide and ethyl 3-iodobenzoate.

20 ¹H NMR/CDCl₃: 1.35 (s, 6H); 1.95 (m, 2H), 3.02 (m, 2H), 6.94 (t, 1H), 7.18 (dd, 1H); 7.33 to 7.39 (m, 2H), 7.61 (dd, 1H), 8.08 (dd, 1H), 8.16 (d, 1H).

EXAMPLE 12:

25 6-(4,4-Dimethylthiochroman-8-ylselenalyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 4,4-dimethylthiochroman-8-diselenide and ethyl 6-iodonicotinate.

^1H NMR/ CDCl_3 : 1.37 (s, 6H); 1.95 (m, 2H), 2.97 (m, 2H), 6.90 (d, 1H), 7.04 (t, 1H); 7.48 to 7.57 (m, 2H), 7.96 (dd, 1H), 9.03 (d, 1H).

EXAMPLE 13:

5 4-(5,5,8,8-Tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)benzoic acid

(a) 5,6,7,8-Tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

A 1.7 M solution of tert-butyllithium in
 10 pentane (37.4 mmol, 22 ml) is added to a solution of 2-bromo-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-naphthalene (4.22 g, 15.8 mmol) in THF (100 ml) at -78°C over 10 min. The mixture is stirred at 0°C for 30 min. Selenium (1.33 g, 16.8 mmol) is added in
 15 2 portions. The mixture is stirred at 0°C for 15 min and then at room temperature for 30 min. 1N HCl solution (40 ml) is added and the reaction mixture is then treated with ethyl ether. The organic phase is washed twice with water, dried over anhydrous magnesium
 20 sulphate and concentrated on a rotary evaporator under vacuum at 40°C. 10 ml of ethanol and 50 mg of sodium hydroxide are added to the oil obtained. The mixture is stirred vigorously for a few minutes in air (until all the product has precipitated) and is then concentrated
 25 on a rotary evaporator under vacuum at 40°C. The solid obtained is filtered off on silica (eluting with heptane) and is then crystallized from an ethanol/ether mixture.

091919-0320
 108220-67267260

Orange solid. Mass: 2.9 g. Yield: 69%.

^1H NMR (CDCl_3): 1.21 (6H, s), 1.25 (6H, s), 1.65 (4H, s), 7.20 (1H Ar, d, $J=8.25$ Hz), 7.38 (1H Ar, dd, $J=1.9$ Hz, $J=8.25$ Hz), 7.51 (1H Ar, d, $J=1.9$ Hz).

- 5 (b) 4-(5,5,8,8-Tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenalyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl
10 4-iodobenzoate.

^1H NMR/ CDCl_3 : 1.26 (s, 6H); 1.30 (s, 6H), 1.70 (s, 4H), 7.27 to 7.37 (m, 4H), 7.54 (d, 1H), 7.91 (d, 2H).

EXAMPLE 14:

- 15 3-(5,5,8,8-Tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenalyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodobenzoate.

- 20 ^1H NMR/ CDCl_3 : 1.25 (s, 6H); 1.27 (s, 6H), 1.68 (s, 4H), 7.24 to 7.26 (m, 2H), 7.34 (t, 1H), 7.48 (s, 1H), 7.60 (dd, 1H), 7.94 (dd, 1H), 8.19 (d, 1H).

EXAMPLE 15:

- 25 6-(5,5,8,8-Tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenalyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl

103250-6267-60

6-iodonicotinate.

^1H NMR/ CDCl_3 : 1.29 (s, 6H); 1.32 (s, 6H), 1.72 (s, 4H), 7.03 (s, 1H), 7.36 (d, 1H), 7.45 (dd, 1H), 7.65 (d, 1H), 7.99 (dd, 1H), 9.07 (d, 1H).

5 **EXAMPLE 16:**

4-[5-Adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenylselenalyl]benzoic acid

a) 5-Adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenyl diselenide

10 A small portion of a solution of
2-(adamantan-1-yl)-4-bromo-5-methyl-
1-methoxyethoxymethoxyphenyl (17 g, 41.5 mmol) in THF
(160 ml) is poured onto a mixture of magnesium (1.51 g)
and one crystal of iodine, with gentle heating. When
15 the reaction medium decolourizes, the rest of the
solution is added so as to maintain a gentle reflux.
After the end of the addition, the solution is refluxed
for 1 h. After cooling to room temperature, 3.6 g of
selenium are added. The reaction medium is stirred for
20 3 h at room temperature and 1N hydrochloric acid
solution (105 ml) and ethyl ether are then added to the
reaction medium. The organic phase is washed with
water, dried over magnesium sulphate and concentrated
on a rotary evaporator. Sodium hydroxide (131 mg) and
25 ethanol (27 ml) are then added. The suspension is
stirred in air and at room temperature for 12 h. The
product is purified by filtration on silica, eluting
with dichloromethane. 12 g (71%) of a yellow solid are

obtained. m.p. = 101°C.

¹H NMR/CDC13: 1.73 (s, 6H); 2.00 (s, 9H); 2.30 (s, 3H); 3.40 (s, 3H); 3.59 (m, 2H); 3.83 (m, 2H); 5.29 (s, 2H); 6.95 (s, 1H); 7.48 (s, 1H).

- 5 b) 4-[5-Adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenylselenanyl]benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 5-adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenyl diselenide and ethyl 4-iodobenzoate.

¹H NMR/CDC13: 1.75 (s, 6H); 2.07 (s, 9H), 2.34 (s, 3H), 3.42 (s, 3H), 3.62 (m, 2H), 3.89 (m, 2H), 5.35 (s, 2H), 7.14 (s, 1H), 7.19 (d, 2H), 7.50 (s, 1H), 7.87 (d, 2H).

EXAMPLE 17:

- 15 3-[5-Adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenylselenanyl]benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 5-adamantan-1-yl-4-(2-methoxyethoxymethoxy)-2-methylphenyl diselenide and ethyl 3-iodobenzoate.

¹H NMR/CDC13: 1.75 (s, 6H); 2.06 (s, 9H), 2.34 (s, 3H), 3.41 (s, 3H), 3.62 (m, 2H), 3.87 (m, 2H), 5.34 (s, 2H), 7.10 (s, 1H), 7.28 (t, 1H), 7.38 (dd, 1H), 7.47 (s, 1H), 7.87 (dd, 1H), 8.02 (d, 1H).

- 25 EXAMPLE 18:

6-(4-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)nicotinic acid

- a) 4-Methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-

09719219-032804
T08220-0426260

tetramethylnaphthalene-2-diselenide

In a manner similar to that of Example 1(a), starting with 2-bromo-5,5,8,8-tetramethyl-4-methoxyethoxymethoxy-5,6,7,8-tetrahydronaphthalene, the expected compound is obtained in the form of an orange oil.

b) 6-(4-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 4-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 6-iodonicotinate.

^1H NMR/ CDCl_3 : 1.27 (s, 6H); 1.42 (s, 6H), 1.67 (m, 4H), 3.36 (s, 3H), 3.56 (m, 2H), 3.82 (m, 2H), 5.29 (s, 2H), 7.11 (d, 1H), 7.31 (d, 1H), 7.35 (d, 1H), 8.00 (dd, 1H), 9.06 (d, 1H).

EXAMPLE 19:

3-(4-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 4-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 3-iodobenzoate.

^1H NMR/ CDCl_3 : 1.26 (s, 6H); 1.38 (s, 6H), 1.62 (m, 4H), 3.36 (s, 3H), 3.53 (m, 2H), 3.78 (m, 2H), 5.22 (s, 2H), 7.12 (d, 1H), 7.15 (d, 1H), 7.35 (t, 1H), 7.65 (dd,

0919219-03201
T0320-6T26T/60

1H), 7.96 (dd, 1H), 8.20 (d, 1H).

EXAMPLE 20:

4-(4-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)-3-methoxybenzoic acid

5 The product is obtained in a manner similar to that of Example 7, starting with 4-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodo-3-methoxybenzoate.

10 ¹H NMR/CDCl₃: 1.26 (s, 6H); 1.42 (s, 6H), 1.66 (m, 4H), 3.35 (s, 3H), 3.54 (m, 2H), 3.81 (m, 2H), 3.98 (s, 3H), 5.27 (s, 2H), 6.94 (d, 1H), 7.25 (d, 1H), 7.30 (d, 1H), 7.48 to 7.53 (m, 2H).

EXAMPLE 21:

15 **3-(4-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)-4-methoxybenzoic acid**

 The product is obtained in a manner similar to that of Example 7, starting with 4-methoxyethoxymethoxy-5,6,7,8-tetrahydro-20 5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 3-iodo-4-methoxybenzoate.

¹H NMR/CDCl₃: 1.25 (s, 6H); 1.40 (s, 6H), 1.65 (m, 4H), 3.34 (s, 3H), 3.53 (m, 2H), 3.80 (m, 2H), 3.97 (s, 3H), 5.26 (s, 2H), 6.88 (d, 1H), 7.21 (d, 1H), 7.24 (d, 1H), 25 7.82 (d, 1H), 7.94 (dd, 1H).

09719219-032801

EXAMPLE 22:

6-(4-Methoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

- a) 4-Methoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

In a manner similar to that of Example 1(a), starting with 2-bromo-5,5,8,8-tetramethyl-4-methoxymethoxy-5,6,7,8-tetrahydronaphthalene, the expected compound is obtained in the form of an orange oil.

- b) 6-(4-Methoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 4-methoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 6-iodonicotinate.

^1H NMR/ CDCl_3 : 1.27 (s, 6H); 1.43 (s, 6H), 1.67 (m, 4H), 3.49 (s, 3H), 5.20 (s, 2H), 7.11 (d, 1H), 7.24 (d, 1H), 7.35 (d, 1H), 8.01 (dd, 1H), 9.07 (d, 1H).

EXAMPLE 23:

6-(3-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

- a) 3-Methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

In manner similar to that of Example 1(a), starting with 2-bromo-5,5,8,8-tetramethyl-3-methoxyethoxymethoxy-5,6,7,8-tetrahydronaphthalene, the expected compound is obtained in the form of an

09494-0320
T08220-6T26T260

orange oil.

b) 6-(3-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 6-iodonicotinate.

¹H NMR/CDCl₃: 1.25 (s, 6H); 1.31 (s, 6H), 1.69 (s, 4H), 3.36 (s, 3H), 3.51 (m, 2H), 3.74 (m, 2H), 5.22 (s, 2H), 7.04 (d, 1H), 7.23 (s, 1H), 7.61 (s, 1H), 7.97 (dd, 1H), 9.05 (d, 1H).

EXAMPLE 24:

2-(3-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 2-iodonicotinate.

¹H NMR/CDCl₃: 1.25 (s, 6H); 1.31 (s, 6H), 1.68 (s, 4H), 3.37 (s, 3H), 3.52 (m, 2H), 3.74 (m, 2H), 5.17 (s, 2H), 7.10 (dd, 1H), 7.22 (s, 1H), 7.54 (s, 1H), 8.29 (dd, 1H), 8.44 (dd, 1H).

25 EXAMPLE 25:

4-(3-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid

The product is obtained in a manner similar

09719219-032801
TOPSECRET

to that of Example 7, starting with
3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-
5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl
4-iodobenzoate.

- 5 ^1H NMR/ CDCl_3 : 1.19 (s, 6H); 1.29 (s, 6H), 1.63 (s, 4H),
3.36 (s, 3H), 3.50 (m, 2H), 3.71 (m, 2H), 5.22 (s, 2H),
7.16 (s, 1H), 7.36 (s, 1H), 7.41 (d, 2H), 7.93 (d, 2H).

EXAMPLE 26:

- 3-(3-Methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-
10 tetrahydro-2-naphthylselanyl)benzoic acid

The product is obtained in a manner similar
to that of Example 7, starting with
3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-
5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl

- 15 3-iodobenzoate.

^1H NMR/ CDCl_3 : 1.12 (s, 6H); 1.27 (s, 6H), 1.63 (m, 4H),
3.37 (s, 3H), 3.52 (m, 2H), 3.77 (m, 2H), 5.26 (s, 2H),
7.12 (s, 1H), 7.13 (s, 1H), 7.38 (t, 1H), 7.69 (dd,
1H), 7.99 (dd, 2H), 8.25 (d, 1H).

- 20 **EXAMPLE 27:**

6-(3,5-Di-tert-butyl-2-methoxymethoxyphenylselanyl)-
nicotinic acid

- a) 2-Bromo-4,6-di-tert-butyl-1-methoxymethoxyphenyl.

A mixture of 2-bromo-4,6-di-tert-butylphenol
25 (4.4 mmol), caesium carbonate (2.95 g) and
methoxymethyl chloride (4.8 mmol) in DMF (18 ml) is
stirred at room temperature for 24 h. The reaction
medium is extracted with ethyl ether. The organic phase

is washed with water, dried over magnesium sulphate and concentrated on a rotary evaporator. The product is purified by filtration on silica.

- b) 4,6-Di-tert-butyl-1-methoxymethoxyphen-2-yl
5 diselenide.

In a manner similar to that of Example 10(c), starting with 10 g of the product obtained above, 1.1 g of magnesium and 2.63 g of selenium, 7.6 g (76%) of the expected product are obtained in the form of a yellow
10 solid.

¹H NMR/CDCl₃: 1.18 (s, 9H); 1.42 (s, 9H); 3.68 (s, 3H); 5.08 (s, 2H); 7.23 (d, 1H); 7.54 (d, 1H).

- c) 6-(3,5-Di-tert-butyl-2-methoxymethoxyphenyl-selanyl)nicotinic acid
15 The product is obtained in a manner similar to that of Example 7, starting with 4,6-di-tert-butyl-1-methoxymethoxyphen-2-yl diselenide and ethyl 6-iodonicotinate.

¹H NMR/CDCl₃: 1.30 (s, 9H); 1.45 (s, 9H); 3.51 (s, 3H);
20 5.17 (s, 2H); 6.94 (d, 1H); 7.50 (d, 1H), 7.56 (d, 1H), 7.98 (dd, 1H), 9.05 (d, 1H).

EXAMPLE 28:

2-(3,5-Di-tert-butyl-2-methoxymethoxyphenylselanyl)-
nicotinic acid

- 25 The product is obtained in a manner similar to that of Example 7, starting with 4,6-di-tert-butyl-1-methoxymethoxyphen-2-yl diselenide and ethyl 2-iodonicotinate.

¹H NMR/CDCl₃: 1.30 (s, 9H); 1.46 (s, 9H); 3.51 (s, 3H); 5.16 (s, 2H); 7.12 (dd, 1H); 7.44 (s, 1H), 8.30 (dd, 1H), 8.46 (dd, 1H).

EXAMPLE 29:

5 **4-(3,5-Di-tert-butyl-2-methoxymethoxyphenylselanyl)-benzoic acid**

The product is obtained in a manner similar to that of Example 7, starting with 4,6-di-tert-butyl-1-methoxymethoxyphen-2-yl diselenide and ethyl
10 4-iodobenzoate.

¹H NMR/CDCl₃: 1.25 (s, 9H); 1.44 (s, 9H); 3.55 (s, 3H); 5.15 (s, 2H); 7.33 to 7.41 (m, 3H); 7.92 (d, 2H).

EXAMPLE 30:

15 **3-(3,5-Di-tert-butyl-2-methoxymethoxyphenylselanyl)-benzoic acid**

The product is obtained in a manner similar to that of Example 7, starting with 4,6-di-tert-butyl-1-methoxymethoxyphen-2-yl diselenide and ethyl
3-iodobenzoate.

20 ¹H NMR/CDCl₃: 1.20 (s, 9H); 1.44 (s, 9H); 3.60 (s, 3H); 5.17 (s, 2H); 7.15 (d, 1H), 7.32 to 7.36 (m, 2H), 7.56 (dd, 1H); 7.96 (dd, 1H), 8.18 (d, 1H).

EXAMPLE 31:

25 **6-[4-Adamantan-1-yl-3-benzyloxyphenylselenalyl]-nicotinic acid**

a) 2-(Adamantan-1-yl)-5-bromo-1-(2-methoxyethoxymethoxy)phenyl

60% sodium hydride (2.5 g) is added

05719219-032801
TABLED " 67267250

portionwise to a solution of 2-(adamantan-1-yl)-5-bromo-1-phenol (20.9 g) in a mixture of THF and DMF (5/5). Stirring is continued for 30 min at room temperature after the end of the addition, and

5 methoxyethoxymethyl chloride (8.92 g) is then added.

The reaction medium is stirred for 4 h at room temperature and is then treated with water and ethyl ether. The organic phase is washed with water, dried over magnesium sulphate and concentrated. After

10 filtration on silica, 17 g (64%) of the expected product are obtained in the form of a white solid.
m.p. = 88°C.

b) 4-Adamantan-1-yl-3-(2-methoxyethoxymethoxy)phenyl diselenide.

15 In a manner similar to that of Example 1(a), starting with 13.04 g of 2-(adamantan-1-yl)-5-bromo-1-methoxyethoxymethoxyphenyl, 9.9 g (76%) of the expected product are obtained in the form of a yellow oil.

20 ¹H NMR/CDC13: 1.55 (s, 6H); 2.05 (d, 9H); 3.38 (s, 3H); 3.57 (m, 2H); 3.82 (m, 2H); 5.27 (s, 2H), 7.11 (d, 1H); 7.22 (dd, 1H); 7.38 (d, 1H).

c) 4-Adamantan-1-yl-3-hydroxyphenyl diselenide

A mixture of the product obtained above
25 (200 mg), concentrated sulphuric acid (1.4 ml), methanol (20 ml) and THF (20 ml) is stirred for 12 h at room temperature. The reaction medium is extracted with ethyl acetate. The organic phase is washed twice with

water, dried over magnesium sulphate and concentrated on a rotary evaporator under vacuum. The expected product is purified by flash chromatography to give an orange-coloured powder.

5 d) 4-Adamantan-1-yl-3-benzyloxyphenyl diselenide.

A mixture of the product obtained above (4.4 mmol), caesium carbonate (2.95 g) and benzyl chloride (1.3 ml) in DMF (18 ml) is stirred at room temperature for 24 h. The reaction medium is extracted
10 with ethyl ether. The organic phase is washed with water, dried over magnesium sulphate and concentrated on a rotary evaporator. The product is purified by filtration on silica (heptane and then dichloromethane). The expected compound is obtained in
15 the form of a yellow powder.

e) 6-[4-Adamantan-1-yl-3-benzyloxyphenylselenanyl]-nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 4-adamantan-1-yl-
20 3-benzyloxyphenyl diselenide and ethyl 6-iodonicotinate.

^1H NMR/ CDCl_3 , acetone D_6 : 1.74 (s, 6H); 2.06 (s, 3H); 2.17 (s, 6H); 5.12 (s, 2H); 6.97 (d, 1H), 7.26 to 7.48 (m, 8H), 7.95 (dd, 1H), 9.04 (d, 1H).

25 **EXAMPLE 32:**

6-(3,5-Di-tert-butyl-2-benzyloxyphenylselenanyl)nicotinic acid

a) 3,5-Di-tert-butyl-2-benzyloxyphenyl diselenide

The procedure is identical to that followed for Example 31(c) and 31(d), applied to the product of Example 27(b).

b) 6-(3,5-Di-tert-butyl-2-benzyloxyphenylselanyl)-
5 nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 3,5-di-tert-butyl-2-benzyloxyphenyl diselenide and ethyl 6-iodonicotinate.

10 $^1\text{H}/\text{CDCl}_3$, acetone D_6 : 1.33 (s, 9H); 1.44 (s, 9H); 5.13 (s, 2H); 7.00 (d, 1H); 7.24 to 7.32 (m, 5H), 7.51 (d, 1H), 7.60 (d, 1H), 7.98 (dd, 1H), 9.01 (d, 1H).

EXAMPLE 33:

3-Methoxy-4-(4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-
15 tetramethyl-2-naphthylselanyl)benzoic acid

a) 4-Hydroxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

The product of Example 22(a),
4-methoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-
20 tetramethylnaphthalene-2-diselenide (12.4 g), is treated in a manner similar to that of Example 15(b) to give 11 g (100%) of the expected compound in the form of a yellow solid. m.p. = 200°C.

^1H NMR/ CDCl_3 : 1.22 (s, 6H); 1.42 (s, 6H); 1.63 (m, 4H);
25 5.25 (s, 1H); 6.75 (d, 1H); 7.11 (d, 1H)

b) 4-Benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

A mixture of the product obtained above

09719219-032801
FOUOED-6726760

(2.5 g, 4.4 mmol), caesium carbonate (2.95 g) and benzyl chloride (1.3 ml) in DMF (18 ml) is stirred at room temperature for 24 h. The reaction medium is extracted with ethyl ether. The organic phase is washed with water, dried over magnesium sulphate and concentrated on a rotary evaporator. The product is purified by filtration on silica (heptane and then dichloromethane). 2.1 g (63%) of the expected compound are obtained in the form of a yellow powder.

10 ^1H NMR/ CDCl_3 : 1.21 (s, 6H); 1.34 (s, 6H); 1.59 (m, 4H); 4.96 (s, 2H); 7.02 (d, 1H); 7.21 (d, 1H); 7.29 to 7.41 (m, 5H).

c) 3-Methoxy-4-(4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselenanyl)benzoic acid

15 The product is obtained in a manner similar to that of Example 7, starting with 4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodo-3-methoxybenzoate.

^1H NMR/ CDCl_3 : 1.27 (s, 6H), 1.43 (s, 6H), 1.66 (m, 4H), 20 3.98 (s, 3H), 5.04 (s, 2H), 6.88 (d, 1H), 7.01 (d, 1H), 7.29 (s, 1H), 7.33 to 7.52 (m, 7H).

EXAMPLE 34:

4-(4-Benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselenanyl)benzoic acid

25 The product is obtained in a manner similar to that of Example 7, starting with 4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodobenzoate.

09719219-032204

¹H NMR/CDCl₃: 1.26 (s, 6H), 1.41 (s, 6H), 1.65 (m, 4H), 5.02 (s, 2H), 6.92 (d, 1H), 7.22 (d, 1H), 7.31 to 7.41 (m, 7H), 7.90 (d, 2H).

EXAMPLE 35:

- 5 **6-(4-Benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl)nicotinic acid**

The product is obtained in a manner similar to that of Example 7, starting with 4-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-

- 10 2-diselenide and ethyl 6-iodonicotinate.

¹H NMR/CDCl₃: 1.28 (s, 6H), 1.43 (s, 6H), 1.67 (m, 4H), 5.07 (s, 2H), 7.00 (d, 1H), 7.04 (d, 1H), 7.32 to 7.44 (m, 6H), 7.96 (dd, 1H), 9.06 (d, 1H).

EXAMPLE 36:

- 15 **3-Methoxy-4-(3-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl)benzoic acid**

a) 3-Hydroxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

The product of Example 23(a),

- 20 3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide, is treated in a manner similar to that of Example 31(c) to give the expected compound in the form of a yellow solid (100%).

- b) 3-Benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide
- 25

The product above is treated in a manner similar to that of Example 33(b).

c) 3-Methoxy-4-(3-benzyloxy-5,6,7,8-tetrahydro-

09719219-032301
TOP SECRET/50

5,5,8,8-tetramethyl-2-naphthylselenanyl)benzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodo-3-methoxybenzoate.

¹H NMR/CDCl₃: 1.22 (s, 6H), 1.25 (s, 6H), 1.67 (s, 4H), 3.97 (s, 3H), 5.07 (s, 2H), 6.89 (d, 1H), 6.90 (d, 1H), 7.22 to 7.25 (m, 5H), 7.50 to 7.53 (m, 3H).

EXAMPLE 37:

6-(3-Benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselenanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-benzyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 6-iodonicotinate

¹H NMR/acetone D₆, CDCl₃: 1.25 (s, 6H), 1.27 (s, 6H), 1.68 (s, 4H), 5.08 (s, 2H), 6.94 (s, 1H), 7.04 (d, 1H), 7.31 (s, 3H), 7.62 (s, 1H), 7.94 (dd, 1H), 9.04 (d, 1H).

EXAMPLE 38:

4-(3-Hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselenanyl)-3-methoxybenzoic acid

a) 3-Hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthylene-2-diselenide

60% sodium hydride (225 mg, 5.63 mmol) is added portionwise to a solution of 4-hydroxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide (1.2 g, 2.56 mmol) in 15 ml of THF and 15 ml of THF.

Stirring is continued for 30 min at room temperature after the end of the addition, and iodohexane (1 ml, 6.8 mmol) is then added. The reaction medium is stirred for 4 h at room temperature and is then treated with water and ethyl ether. The organic phase is washed with water, dried over magnesium sulphate and concentrated. After purification by chromatography on silica (95 heptane/5 CH₂Cl₂), the product is obtained in the form of a yellow oil.

¹H NMR/CDCl₃: 0.90 (m, 9H); 1.30 to 1.48 (m, 12H); 1.59 (m, 4H); 1.77 (m, 2H); 3.85 (t, 2H); 6.92 (d, 1H); 7.17 (d, 1H).

b) 4-(3-Hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl)-3-methoxybenzoic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide and ethyl 4-iodo-3-methoxybenzoate.

¹H NMR/CDCl₃: 0.89 (t, 3H), 1.27 (s, 6H), 1.30 to 1.37 (m, 4H), 1.42 (s, 6H), 1.48 (m, 2H), 1.63 (m, 4H), 1.82 (m, 2H), 3.90 (t, 2H), 3.98 (s, 3H), 6.91 (d, 1H), 6.93 (s, 1H), 7.24 (s, 1H), 7.49 to 7.55 (m, 2H).

EXAMPLE 39:

6-(3-Hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl)nicotinic acid

The product is obtained in a manner similar to that of Example 7, starting with 3-hexyloxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide

and ethyl 6-iodonicotinate.

¹H NMR/CDCl₃: 0.89 (t, 3H), 1.27 (s, 6H), 1.30 to 1.37 (m, 4H), 1.42 (s, 6H), 1.48 (m, 2H), 1.63 (m, 4H), 1.84 (m, 2H), 3.92 (t, 2H), 6.97 (d, 1H), 7.08 (d, 1H), 7.29 (d, 1H), 8.00 (dd, 1H), 9.08 (d, 1H).

EXAMPLE 40:

4-(5-Adamantan-1-yl-4-benzyloxy-2-methylphenyl-selenalyl)benzoic acid

a) 5-Adamantan-1-yl-4-benzyloxy-2-methylphenyl
10 diselenide

The procedure is identical to that followed for Example 31(c) and 31(d), applied to the product of Example 16(a).

b) 4-(5-Adamantan-1-yl-4-benzyloxy-2-methylphenyl-selenalyl)benzoic acid
15

The product is obtained in a manner similar to that of Example 7, starting with 5-adamantan-1-yl-4-benzyloxy-2-methylphenyl diselenide and ethyl 4-iodobenzoate.

20 ¹H NMR/acetone D₆, CDCl₃: 1.70 (s, 6H); 2.02 (s, 3H), 2.11 (s, 6H), 2.41 (s, 3H), 5.16 (s, 2H), 6.85 (dd, 1H), 6.98 (s, 1H), 7.35 to 7.58 (m, 6H), 7.97 (dd, 2H), 9.05 (d, 1H).

EXAMPLE 41:

25 Ethyl 6-[3-[5-(tert-butyldimethylsilanyloxy)pentyloxy-methyl]-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl)nicotinate

a) 5-(3-Bromo-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-

09719249-032804
FORCED "67267/60

2-naphthyloxy)pentyl acetate

A solution of 3-bromo-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-ol (10 g, 0.35 mol), 5-bromopentyl acetate (8.15 g) and potassium carbonate (33.6 g) in methyl ethyl ketone (200 ml) is refluxed for 2 hours. The reaction medium is treated with water and ethyl acetate. After separation of the phases by settling, the organic phase is washed twice with water, dried over magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C. The product is purified by flash chromatography on a column of silica. Yellow oil. Yield: 93%.

b) [5-(3-Bromo-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthyloxy)pentyl]-tert-butyltrimethylsilane

The acetate obtained above is saponified and the resulting hydroxyl group is then protected according to the following procedure: tert-butyltrimethylsilyl chloride (2.64 g) is added to a mixture of 5-(3-bromo-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthyloxy)pentan-1-ol (4.3 g, 11.7 mmol) and 80% sodium hydride (422 mg) in THF (20 ml).

The mixture is stirred at room temperature for 2 h. The solution is poured into a mixture of water and ethyl acetate. The organic phase is washed twice with water, dried over magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C. The product is purified by flash chromatography

09719219-032801

on a column of silica.

Yellow oil. Yield: 64%.

- c) 3-[5-(*tert*-Butyldimethylsilanyloxy)pentyloxy]-
5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalene-
5 2-diselenide

The expected product is obtained from the
bromo derivative obtained above, in a manner similar to
that of Example 1a. Yellow oil. Yield: 10%.

- d) Ethyl 6-[3-[5-(*tert*-butyldimethylsilanyloxy)-
10 pentyloxy]-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
2-naphthylselanyl]nicotinate.

In a manner similar to that of Example 1(b),
by reaction of 257 mg (0.27 mmol) of the diselenide
obtained above in 25 ml of ethanol with 119 mg of
15 sodium borohydride, 120 mg (0.43 mmol) of ethyl
6-iodonicotinate and 4 mg of bis(bipyridine)nickel(II)
dibromide, 152 mg (56%) of the expected derivative are
obtained in the form of a yellow oil.

- ¹H NMR (CDCl₃): 0.00 (6H, s), 0.85 (9H, s), 1.22
20 (6H, s), 1.30 (6H, s), 1.33 to 1.50 (6H, m), 1.60 to
1.67 (7H, m), 3.48 (2H, t), 3.92 (2H, t), 4.35 (2H, q),
6.84 (1H, s), 6.99 (1H, d), 7.57 (1H, s), 7.91
(1H, dd), 8.97 (1H, d).

EXAMPLE 42:

- 25 6-[3-[5-(*tert*-Butyldimethylsilanyloxy)pentyloxy]-
5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-
2-naphthylselanyl]nicotinic acid

In a manner similar to that of Example 2, by

5

EXAMPLE 43:

6-[3-(5-Hydroxypentyloxy)-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthylselanyl]nicotinic acid

10

25 **EXAMPLE 44:**

Ethyl 4-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate

(a) 5,6,7,8-Tetrahydro-5,5,8,8-tetramethylnaphthalene-

2-diselenide

A 1.7M solution of tert-butyllithium in pentane (37.4 mmol, 22 ml) is added to a solution of 2-bromo-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-naphthalene (4.22 g, 15.8 mmol) in THF (100 ml) at -78°C over 10 min. The mixture is stirred at 0°C for 30 min. Selenium (1.33 g, 16.8 mmol) is added in 2 portions. The mixture is stirred at 0°C for 15 min and then at room temperature for 30 min. 1N HCl solution (40 ml) is added and the reaction mixture is then treated with ethyl ether. The organic phase is washed twice with water, dried over anhydrous magnesium sulphate and concentrated on a rotary evaporator under vacuum at 40°C. 10 ml of ethanol and 50 mg of sodium hydroxide are added to the oil obtained. The mixture is stirred vigorously for a few minutes in air (until all the product has precipitated) and is then concentrated on a rotary evaporator under vacuum at 40°C. The solid obtained is filtered off on silica (eluting with heptane) and then crystallized from an ethanol/ether mixture.

Orange solid. Mass: 2.9 g. Yield: 69%.

¹H NMR (CDCl₃): 1.21 (6H, s), 1.25 (6H, s), 1.65 (4H, s), 7.20 (1H Ar, d, J=8.25 Hz), 7.38 (1H Ar, dd, J=1.9 Hz, J=8.25 Hz), 7.51 (1H Ar, d, J=1.9 Hz).

b) Ethyl 4-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)benzoate

In a manner similar to that of Example 1(b),

by reaction of 213 mg (0.4 mmol) of the diselenide obtained above in 20 ml of ethanol with 73 mg of sodium borohydride (1.92 mmol), 177 mg (0.64 mmol) of ethyl 4-iodobenzoate and 37 mg of tetrakis(triphenyl-

5 phosphine)palladium, and after purification by flash chromatography (70 heptane/30 CH₂Cl₂), 151 mg of the expected derivative are obtained in the form of a yellow solid. m.p. = 73°C.

1H NMR (CDCl₃): 1.26 (6H, s), 1.29 (6H, s), 1.37 (t, 3H), 1.70 (4H, s), 4.34 (q, 2H), 7.15 to 7.25 (m, 3H), 7.32 (1H, d), 7.44 (1H, d), 7.89 (1H, d).

10

EXAMPLE 45:

Ethyl 4-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselenanyl)benzoate

15 In a manner similar to that of Example 1(b), by reaction of 3.35 g (4.5 mmol) of 3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-tetramethylnaphthalene-2-diselenide in 100 ml of ethanol with 501 mg of sodium borohydride (13.5 mmol),

20 2.5 g (9 mmol) of ethyl 4-iodobenzoate and 90 mg of bis(bipyridine)nickel(II) dibromide, and after purification by flash chromatography (85 heptane/15 EtOAc), 2.58 g of the expected derivative are obtained in the form of a yellow oil

25 (83%).

EXAMPLE 46:

Ethyl 4-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate

A mixture of ethyl 4-(3-methoxyethoxymethoxy-
5 5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
2-naphthylselanyl)benzoate (2.3 g, 4.4 mmol),
concentrated sulphuric acid (475 μ l), methanol (40 ml)
and THF (20 ml) is stirred for 48 h at room temperature.
The reaction medium is extracted with ethyl ether. The
10 organic phase is washed twice with water, dried over
magnesium sulphate and concentrated on a rotary
evaporator under vacuum. The product is purified by
crystallization from heptane. 2.06 g (97%) of the
expected compound are obtained in the form of an
15 orange-coloured powder. m.p. = 113°C.

EXAMPLE 47:

4-(3-Hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid

In a manner similar to that of Example 2, by
20 reaction of 400 mg (0.92 mmol) of ethyl 4-(3-hydroxy-
5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
2-naphthylselanyl)benzoate with 336 mg (8.4 mmol) of
sodium hydroxide in a THF/ethanol mixture
(20 ml/20 ml), 214 mg (58%) of a pink powder are
25 obtained. m.p. = 217°C.

0919219-032804

EXAMPLE 48:

Ethyl 6-(3-methoxyethoxymethoxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate

In a manner similar to that of Example 1(b),
 5 by reaction of 3.35 g (4.5 mmol) of
 3-methoxyethoxymethoxy-5,6,7,8-tetrahydro-5,5,8,8-
 tetramethylnaphthalene-2-diselenide in 100 ml of
 ethanol with 501 mg of sodium borohydride (13.5 mmol),
 2.5 g (9 mmol) of ethyl 4-iodobenzoate and 90 mg of
 10 bis(bipyridine)nickel(II) dibromide, and after
 purification by flash chromatography
 (85 heptane/15 EtOAc), 2.09 g of the expected
 derivative are obtained in the form of a yellow oil
 (45%).

15 ¹H NMR/CDCl₃: 1.25 (s, 6H), 1.31 (s, 6H), 1.38 (t, 3H),
 1.69 (m, 4H), 3.36 (s, 3H), 3.50 (m, 2H), 3.73 (m, 2H),
 4.37 (q, 2H), 5.22 (s, 2H), 7.01 (d, 1H), 7.22 (s, 1H),
 7.60 (s, 1H), 7.94 (dd, 1H), 8.99 (d, 1H).

EXAMPLE 49:

20 **Ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-
 tetrahydro-2-naphthylselanyl)nicotinate**

A mixture of ethyl 6-(3-methoxyethoxymethoxy-
 5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)nicotinate (2.6 g, 5 mmol),
 25 concentrated sulphuric acid (535 μ l), ethanol (75 ml)
 and THF (25 ml) is stirred for 3 days at room
 temperature. The reaction medium is extracted with
 ethyl ether. The organic phase is washed twice with

09719219-032801

water, dried over magnesium sulphate and concentrated on a rotary evaporator under vacuum. The solid obtained is washed with ethyl ether. 2.01 g (93%) of the expected compound are obtained in the form of an
 5 orange-coloured powder. m.p. = 138°C.

EXAMPLE 50:

6-(3-Hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid

In a manner similar to that of Example 2, by
 10 reaction of 400 mg (0.92 mmol) of ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate with 357 mg (8.9 mmol) of sodium hydroxide in a THF/ethanol mixture (20 ml/20 ml), 60 mg (16%) of a yellow powder are
 15 obtained. m.p.: 250°C.

EXAMPLE 51:

Ethyl 6-[3-(3-ethoxycarbonylpropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]-nicotinate

20 432 mg (102.0 mmol) of ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate, 276 mg (2 mmol) of potassium carbonate and 390 mg (2 mmol) of ethyl 4-bromobutanoate are introduced into a three-necked
 25 flask. The mixture is heated at 80°C for 12 h. The reaction medium is poured into water and extracted with ethyl ether, and the organic phase is separated out by settling, washed with water, dried over magnesium

sulphate and evaporated. After purification by flash chromatography (9 heptane/1 EtOAc), 467 mg (85%) of the expected compound are collected in the form of an orange-coloured oil.

- 5 ^1H NMR/ CDCl_3 : 1.20 to 1.31 (m, 15H), 1.38 (t, 3H), 1.69 (s, 4H), 1.96 (m, 2H), 2.38 (t, 2H), 2.85 (t, 2H), 4.12 (q, 2H), 4.36 (q, 2H), 6.88 (d, 1H), 7.02 (s, 1H), 7.48 (s, 1H), 8.26 (dd, 2H), 8.83 (d, 1H).

EXAMPLE 52:

- 10 **6-[3-(3-Carboxypropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic acid**

- In a manner similar to that of Example 2, by reaction of 340 mg (0.62 mmol) of ethyl 6-[3-(3-ethoxycarbonylpropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate with 250 mg (62.2 mmol) of sodium hydroxide in ethanol (10 ml), 211 mg (69%) of a white powder are obtained. m.p.: 177°C.

EXAMPLE 53:

- 20 **Ethyl 4-[3-(3-ethoxycarbonylpropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]-benzoate**

- In a manner similar to that of Example 51, by reaction of 300 mg (0.86 mmol) of ethyl 4-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate with 336 mg (1.72 mmol) of ethyl 4-bromobutanoate and 238 mg of potassium carbonate in MEK (10 ml), 364 mg (78%) of a yellow oil are obtained.

¹H NMR/CDCl₃: 1.16 to 1.32 (m, 15H), 1.38 (t, 3H), 1.66 (m, 4H), 1.98 (m, 2H), 2.30 (t, 2H), 3.98 (t, 2H), 4.08 (q, 2H), 4.35 (q, 2H), 6.78 (s, 1H), 7.28 (s, 1H), 7.41 (dd, 2H), 7.87 (dd, 2H).

5 **EXAMPLE 54:**

4-[3-(3-Carboxypropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]benzoic acid

In manner similar to that of Example 2, by reaction of 250 mg (0.46 mmol) of ethyl

10 4-[3-(3-carboxypropoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]benzoate with 183 mg (4.6 mmol) of sodium hydroxide in a THF/ethanol mixture (5 ml/5 ml), 172 mg (76%) of a white powder are obtained. m.p.: 230°C.

15 **EXAMPLE 55:**

Ethyl 4-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]-benzoate

In a manner similar to that of Example 51, by
20 reaction of 370 mg (0.86 mmol) of ethyl 4-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate with 408 mg (1.72 mmol) of methyl 8-bromooctanoate and 238 mg of potassium carbonate in MEK (10 ml), 502 mg (99%) of a yellow oil
25 are obtained.

¹H NMR/CDCl₃: 1.16 (s, 6H), 1.26 to 1.29 (m, 12H), 1.38 (t, 3H), 1.56 to 1.68 (m, 8H), 2.28 (t, 2H), 3.66 (s, 3H), 3.92 (t, 2H), 4.36 (q, 2H), 6.78 (s, 1H), 7.27

(s, 1H), 7.41 (dd, 2H), 7.88 (dd, 2H).

EXAMPLE 56:

4-[3-(7-Carboxyheptyloxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]benzoic acid

5 In a manner similar to that of Example 2, by reaction of 410 mg (0.7 mmol) of ethyl 4-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]benzoate with 280 mg (7 mmol) of sodium hydroxide in a THF/ethanol
10 mixture (5 ml/5 ml), 326 mg (85%) of a white powder are obtained. m.p.: 183°C.

EXAMPLE 57:

**Ethyl 6-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]-
15 nicotinate**

 In a manner similar to that of Example 51, by reaction of 460 mg (1.06 mmol) of ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinate with 515 mg (2.17 mmol) of
20 methyl 8-bromooctanoate and 295 mg of potassium carbonate in MEK (10 ml), 487 mg (78%) of a yellow oil are obtained.

EXAMPLE 58:

**6-[3-(7-Carboxyheptyloxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinic [lacuna]
25**

 In a manner similar to that of Example 2, by reaction of 390 mg (0.66 mmol) of ethyl 6-[3-(7-methoxycarbonylheptyloxy)-5,5,8,8-tetramethyl-

09749219-032801

5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate with 265 mg (6.6 mmol) of sodium hydroxide in a THF/ethanol mixture (5 ml/1 ml), 277 mg (77%) of a white powder are obtained. m.p.: 186°C.

5 **EXAMPLE 59:**

Ethyl 6-(3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate

a) 2-Bromoethyl acetate

Acetic anhydride (11.35 ml, 0.12 mol) is added dropwise to a solution of 2-bromoethanol (12.5 g, 0.1 mol) and DMAP (1.22 g) in 125 ml of dichloromethane. The mixture is stirred at room temperature for 12 h and treated with water and dichloromethane. The organic phase is washed with water, dried over magnesium sulphate, concentrated on a rotary evaporator and purified by distillation. Yellowish liquid (94%).

b) Ethyl 6-(3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate

In a manner similar to that of Example 51, by reaction of 477 mg (1.10 mmol) of ethyl 6-(3-hydroxy-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl]nicotinate with 396 mg (2.2 mmol) of 2-bromoethyl acetate and 304 mg of potassium carbonate in MEK (10 ml), 545 mg (96%) of a yellow oil are obtained.

¹H NMR/CDCl₃: 1.24 (s, 6H), 1.32 (s, 6H), 1.38 (t, 3H), 1.69 (s, 4H), 1.99 (s, 3H), 3.00 (t, 3H), 4.24 (t, 2H),

09719219-032801
T08280-6T28T-60

4.38 (q, 2H), 6.89 (d, 1H), 7.03 (s, 1H), 7.54 (s, 1H),
8.27 (dd, 1H), 8.83 (d, 1H).

EXAMPLE 60:

**6-(3-(2-Hydroxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-
5 tetrahydro-2-naphthylselanyl)nicotinic acid**

In a manner similar to that of Example 2, by
reaction of 419 mg (0.81 mmol) of ethyl
6-[3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-
tetrahydro-2-naphthylselanyl]nicotinate with 320 mg
10 (8 mmol) of sodium hydroxide in a THF/ethanol mixture
(4 ml/4 ml), 273 mg (75%) of a white powder are
obtained. m.p.: 170°C.

EXAMPLE 61:

**Ethyl 4-(3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-
15 5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate**

In a manner similar to that of Example 51, by
reaction of 400 mg (0.93 mmol) of ethyl 4-(3-hydroxy-
5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
2-naphthylselanyl)benzoate with 334 mg (2.2 mmol) of
20 2-bromoethyl acetate and 257 mg of potassium carbonate
in MEK (10 ml), 333 mg (69%) of a yellow oil are
obtained.

¹H NMR/CDCl₃: 1.16 (s, 6H), 1.29 (s, 6H), 1.38 (t, 3H),
1.59 (s, 4H), 1.99 (s, 3H), 4.16 (m, 2H), 4.29 to 4.40
25 (m, 4H), 6.82 (s, 1H), 7.28 (s, 1H), 7.43 (d, 1H), 7.89
(d, 1H).

09719219-032801
TOPSECRET

EXAMPLE 62:

4-(3-(2-Hydroxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)benzoic acid

In a manner similar to that of Example 2, by
5 reaction of 322 mg (0.62 mmol) of ethyl
4-[3-(2-acetoxyethoxy)-5,5,8,8-tetramethyl-5,6,7,8-
tetrahydro-2-naphthylselanyl]benzoate with 250 mg
(6.2 mmol) of sodium hydroxide in a THF/ethanol mixture
(3 ml/3 ml), 226 mg (81%) of a white powder are
10 obtained. m.p.: 197°C.

EXAMPLE 63:

**Ethyl 4-(3-(2-chloroethoxy)-5,5,8,8-tetramethyl-
5,6,7,8-tetrahydro-2-naphthylselanyl)benzoate**

In a manner similar to that of Example 51, by
15 reaction of 431 mg (1 mmol) of ethyl 4-(3-hydroxy-
5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-
2-naphthylselanyl benzoate with 222 mg (1.5 mmol) of
1-bromo-2-chloroethyl [lacuna] and 278 mg of potassium
carbonate in MEK (20 ml), 200 mg (40%) of a yellow oil
20 are obtained.

EXAMPLE 64:

**Ethyl 4-[3-(2-iodoethoxy)-5,5,8,8-tetramethyl-5,6,7,8-
tetrahydro-2-naphthylselanyl]benzoate**

A mixture of 200 mg (0.4 mmol) of ethyl
25 4-[3-(2-chloroethoxy)-5,5,8,8-tetramethyl-5,6,7,8-
tetrahydro-2-naphthylselanyl]benzoate with 607 mg
(4 mmol) of sodium iodide in MEK (4 ml) is refluxed for
12 h. The reaction medium is treated with water and

T03220 6 T26T260

ethyl ether. The organic phase is washed with water, dried over magnesium sulphate and concentrated on a rotary evaporator. The oil obtained is reacted under the same conditions. 159 mg (68%) of a yellow solid are
 5 obtained. m.p. = 87°C.

EXAMPLE 65:

6-(3-Adamantan-1-yl-4-methoxyphenylselanyl)nicotinic acid

The product is obtained in a manner similar
 10 to that of Example 7, starting with 3-adamantan-1-yl-4-methoxyphenyl diselenide and ethyl 6-iodonicotinate.
¹H NMR/THF D8: 1.79 (s, 6H), 2.04 (s, 3H), 2.13 (s, 6H), 3.89 (s, 3H), 6.93 (d, 1H), 7.02 (d, 1H), 7.52 to 7.55 (m, 2H), 7.91 (dd, 1H), 8.9 (d, 1H).

15 EXAMPLE 66:

[6-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-3-pyridyl]methanol

3 g (7 mmol) of ethyl 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-
 20 nicotinate, 800 mg (20 mmol) of lithium aluminium hydride and 90 ml of THF are introduced into a round-bottomed flask under a stream of nitrogen. The reaction medium is refluxed for two hours, cooled, the excess hydride is hydrolysed and the salt is filtered off.
 25 After evaporation of the filtrate, the residue obtained is recrystallized from heptane. 1.36 g (50%) of [6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-3-pyridyl]methanol with a melting

09719219-032801

point of 110-111°C are collected.

EXAMPLE 67:

N-Ethyl-6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinamide

- 5 (a) 6-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinyl chloride

2 g (5 mmol) of 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinic acid, 20 ml of toluene, 100 µl of DMF and 450 µl of thionyl
10 chloride are introduced into a round-bottomed flask. The reaction medium is refluxed for one hour and evaporated. 100% of the expected acid chloride are collected, this product being used for the rest of the synthesis without further purification.

- 15 (b) N-Ethyl-6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinamide

By reaction of 2.1 g (5 mmol) of the above acid chloride with 1 ml of ethylamine (70% in water) in 20 ml of THF, 2.02 g (95%) of the expected amide, with
20 a melting point of 218-220°C, are obtained.

EXAMPLE 68:

Morpholin-4-yl-[6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)-3-pyridyl]methanone

By reaction of 2.1 g (5 mmol) of the above
25 acid chloride with 1 ml of morpholine in 20 ml of THF, 2.17 g (93%) of the expected amide, with a melting point of 147-148°C, are obtained.

09749249-032804

EXAMPLE 69:

N-(4-Hydroxyphenyl)-6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthylselanyl)nicotinamide

By reaction of 2.1 g (5 mmol) of the above
 5 acid chloride with 540 mg (5 mmol) of 4-aminophenol in
 40 ml of THF in the presence of 830 μ l of
 triethylamine, 2.35 g (96%) of the expected amide, with
 a melting point of 223-225°C, are obtained.

EXAMPLE 70:

10 **6-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)pyridine-3-carbaldehyde**

By reaction of 890 mg (2.3 mmol) of
 [6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)-3-pyridyl]methanol with 1.12 g
 15 (3 mmol) of pyridinium dichromate in 90 ml of
 dichloromethane, and after filtration on silica, 600 mg
 (68%) of 6-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-
 2-naphthylselanyl)pyridine-3-carbaldehyde, with a
 melting point of 150-152°C, are obtained.

20 **B. FORMULATION EXAMPLES**

1) ORAL ROUTE

(a) The following composition is prepared in the form
 of a 0.8 g tablet

25	Compound of Example 3	0.005 g
	Pregelatinized starch	0.265 g
	Microcrystalline cellulose	0.300 g
	Lactose	0.200 g
	Magnesium stearate	0.030 g

For the treatment of acne, 1 to 3 tablets will be administered to an adult individual per day for 3 to 6 months, depending on the severity of the case treated.

5

(b) A drinkable suspension for packaging in 5 ml vials is prepared:

	Compound of Example 12	0.050 g
	Glycerol	0.500 g
10	70% sorbitol	0.500 g
	Sodium saccharinate	0.010 g
	Methyl parahydroxybenzoate	0.040 g
	Flavouring	qs
	Purified water	qs 5 ml

15

For the treatment of acne, 1 vial will be administered to an adult individual per day for 3 months, depending on the severity of the case treated.

20

(c) The following formulation for packaging in gelatin capsules is prepared:

	Compound of Example 5	0.025 g
	Corn starch	0.060 g
	Lactose qs	0.300 g
25	The gelatin capsules used consist of gelatin, titanium oxide and a preserving agent.	

09719219-03201
T0320-03201

In the treatment of psoriasis, 1 gelatin capsule will be administered to an adult individual per day for 30 days.

2) TOPICAL ROUTE

- 5 (a) The following nonionic water-in-oil cream is prepared:

	Compound of Example 23	0.100 g
	Mixture of emulsifying lanolin	
	alcohols, waxes and refined oils,	
10	sold by the company BDF under the	
	name "anhydrous Eucerin"	39.900 g
	Methyl para-hydroxybenzoate	0.075 g
	Propyl para-hydroxybenzoate	0.075 g
	Sterile demineralized water qs	100.000 g
15	This cream will be applied to psoriatic skin	
	once or twice a day for 30 days.	

- (b) A gel is prepared by making the following formulation:

20	Compound of Example 39	0.050 g
	Base erythromycin	4.000 g
	Butylhydroxytoluene	0.050 g
	Hydroxypropylcellulose sold by	
	the company Hercules under the	
25	name "Kluçel HF"	2.000 g
	Ethanol (at 95°) qs	100.000 g
	This gel will be applied to skin affected	
	with dermatitis or acneic skin 1 to 3 times a day for 6	

1.03220" 67227650

to 12 weeks, depending on the severity of the case treated.

(c) An anti-seborrhoeic lotion is prepared by mixing
5 together the following ingredients:

Compound of Example 6	0.030 g
Propylene glycol	5.000 g
Butylhydroxytoluene	0.100 g
Ethanol (at 95°) qs	100.000 g

10 This lotion will be applied twice a day to a
seborrhoeic scalp, and a significant improvement is
observed within a period of between 2 and 6 weeks.

(d) A cosmetic composition to combat the harmful
15 effects of the sun is prepared by mixing together the
following ingredients:

Compound of Example 59	1.000 g
Benzylidenecamphor	4.000 g
Fatty acid triglycerides	31.000 g
20 Glyceryl monostearate	6.000 g
Stearic acid	2.000 g
Cetyl alcohol	1.200 g
Lanolin	4.000 g
Preserving agents	0.300 g
25 Propylene glycol	2.000 g
Triethanolamine	0.500 g
Fragrance	0.400 g
Demineralized water qs	100.000 g

09719219-032801
TOP SECRET

This composition will be applied daily and makes it possible to combat light-induced ageing.

(e) The following nonionic oil-in-water cream is

5 prepared:

	Compound of Example 16	0.500 g
	Vitamin D3	0.020 g
	Cetyl alcohol	4.000 g
	Glyceryl monostearate	2.500 g
10	PEG-50 stearate	2.500 g
	Karite butter	9.200 g
	Propylene glycol	2.000 g
	Methyl para-hydroxybenzoate	0.075 g
	Propyl para-hydroxybenzoate	0.075 g
15	Sterile demineralized water qs	100.000 g

This cream will be applied to psoriatic skin once or twice a day for 30 days.

(f) A topical gel is prepared by mixing together the
20 following ingredients:

	Compound of Example 4	0.050 g
	Ethanol	43.000 g
	α -tocopherol	0.050 g
	Carboxyvinyl polymer sold under	
25	the name "Carbopol 941" by the	
	company "Goodrich"	0.500 g
	Triethanolamine as an aqueous	
	solution at 20% by weight	3.800 g

05719216-032801
TOP SECRET 67267650

Water 9.300 g

Propylene glycol qs 100.000 g

This gel will be applied in the treatment of
acne 1 to 3 times a day for 6 to 12 weeks, depending on
5 the severity of the case treated.

(g) A hair lotion to combat hair loss and to promote
regrowth of the hair is prepared by mixing together the
following ingredients:

10 Compound of Example 31 0.05 g
Compound sold under the name
"Minoxidil" 1.00 g
Propylene glycol 20.00 g
Ethanol 34.92 g
15 Polyethylene glycol (molecular
mass = 400) 40.00 g
Butylhydroxyanisole 0.01 g
Butylhydroxytoluene 0.02 g
Water qs 100.00 g

20 This lotion will be applied twice a day for
3 months to a scalp which has suffered considerable
hair loss.

(h) An anti-acne cream is prepared by mixing together
25 the following ingredients:

Compound of Example 7 0.050 g
Retinoic acid 0.010 g
Mixture of glycerol stearate and

094949-03301
F08220 6726760

polyethylene glycol stearate (75 mol),
sold under the name "Gelot 64" by
the company "Gattefosse" 15.000 g
Kernel oil polyoxyethylenated with
5 6 mol of ethylene oxide, sold
under the name "Labrafil M2130 CS"
by the company "Gattefosse" 8.000 g
Perhydrosqualene 10.000 g
Preserving agents qs
10 Polyethylene glycol (molecular
mass = 400) 8.000 g
Disodium salt of ethylenediamine-
tetraacetic acid 0.050 g
Purified water qs 100.000 g
15 This cream will be applied to skin affected
with dermatitis or to acneic skin 1 to 3 times a day
for 6 to 12 weeks.

(i) An oil-in-water cream is prepared by making the
20 following formulation:

Compound of Example 43 0.020 g
Betamethasone 17-valerate 0.050 g
S-Carboxymethylcysteine 3.000 g
Polyoxyethylene stearate (40 mol of
25 ethylene oxide) sold under the name
"Myrj 52" by the company "Atlas" 4.000 g
Sorbitan monolaurate, polyoxyethylene
with 20 mol of ethylene oxide, sold

09719219-032801
T08220" 6722T260

under the name "Tween 20" by the
company "Atlas" 1.800 g
Mixture of glyceryl mono- and
distearate sold under the name
5 "Geleol" by the company "Gattefosse". 4.200 g
Propylene glycol 10.000 g
Butylhydroxyanisole 0.010 g
Butylhydroxytoluene 0.020 g
Cetostearyl alcohol 6.200 g
10 Preserving agents qs
Perhydrosqualene 18.000 g
Mixture of caprylic/capric
triglycerides sold under the name
"Miglyol 812" by the company "Dynamit
15 Nobel" 4.000 g
Triethanolamine (99% by weight) 2.500 g
Water qs 100.000 g
This cream will be applied twice a day to
skin affected with dermatitis, for 30 days.

20

(j) The following oil-in-water cream is prepared:

Lactic acid 5.000 g
Compound of Example 1 0.020 g
Polyoxyethylene stearate (40 mol of
25 ethylene oxide) sold under the name
"Myrj 52" by the company "Atlas" 4.000 g
Sorbitan monolaurate, polyoxyethylene
with 20 mol of ethylene oxide, sold

09719219-03801
FOB220" 6T26T/60

under the name "Tween 20" by the
 company "Atlas" 1.800 g
 Mixture of glyceryl mono- and
 distearate sold under the name
 5 "Geleol" by the company "Gattefosse" 4.200 g
 Propylene glycol 10.000 g
 Butylhydroxyanisole 0.010 g
 Butylhydroxytoluene 0.020 g
 Cetostearyl alcohol 6.200 g
 10 Preserving agents qs
 Perhydrosqualene 18.000 g
 Mixture of caprylic/capric
 triglycerides sold under the name
 "Miglyol 812" by the company "Dynamit
 15 Nobel" 4.000 g
 Water qs 100.000 g

This cream will be applied once a day and
 helps to combat ageing, whether this is light-induced
 or chronological ageing.

09719249-032804